


# Frobenius Flocks



Jan Draisma, Universität Bern

j.w.w. Rudi Pendavingh and Guus Bollen, Eindhoven

$K$  a field,  $\sigma$  an automorphism of  $K$ ,  $E$  a finite set  
 $\mathbb{Z}^E$  acts on  $K^E$  via  $\alpha v := (\sigma^{\alpha_i} v_i)_{i \in E}$

$K$  a field,  $\sigma$  an automorphism of  $K$ ,  $E$  a finite set  
 $\mathbb{Z}^E$  acts on  $K^E$  via  $\alpha v := (\sigma^{\alpha_i} v_i)_{i \in E}$

## Definition

$V : \mathbb{Z}^E \rightarrow \{d\text{-dimensional subspaces of } K^E\}, \alpha \mapsto V_\alpha$   
 is a *vector space flock* on  $E$  over  $(K, \sigma)$  of rank  $d$  if

(VF1)  $V_\alpha / i = V_{\alpha + e_i} \setminus i$  and

(VF2)  $V_{\alpha + \mathbf{1}} = \mathbf{1} V_\alpha$ .

$W \setminus i :=$  image of  $W$  in  $K^{E-i}$

$W / i :=$  image of  $e_i^\perp \cap W$  in  $K^{E-i}$

$\mathbf{1}$  = the all-one vector


# Example

3

$$(\text{VF1}) \ V_{\alpha}/i = V_{\alpha+e_i} \setminus i$$

$$(\text{VF2}) \ V_{\alpha+1} = \mathbf{1}V_{\alpha}$$

$$E = \{1, 2\}, \ V_0 = \langle(1, a)\rangle, \ a \neq 0$$



$\langle(1, a)\rangle$

# Example


3

$$(VF1) \ V_{\alpha}/i = V_{\alpha+e_i} \setminus i$$

$$(VF2) \ V_{\alpha+1} = \mathbf{1}V_{\alpha}$$

$$E = \{1, 2\}, \ V_0 = \langle(1, a)\rangle, \ a \neq 0$$

$$V_0/1 = \{0\} = V_{e_1} \setminus 1, \text{ so } V_{e_1} = \langle(1, 0)\rangle$$



# Example


3

$$(VF1) \ V_{\alpha}/i = V_{\alpha+e_i} \setminus i$$

$$(VF2) \ V_{\alpha+1} = \mathbf{1}V_{\alpha}$$

$$E = \{1, 2\}, \ V_0 = \langle(1, a)\rangle, \ a \neq 0$$

$$V_0/1 = \{0\} = V_{e_1} \setminus 1, \text{ so } V_{e_1} = \langle(1, 0)\rangle = V_{ke_1} \text{ for } k \geq 1$$



# Example

3


$$(VF1) V_\alpha / i = V_{\alpha+e_i} \setminus i$$

$$(VF2) V_{\alpha+1} = \mathbf{1}V_\alpha$$

$$E = \{1, 2\}, V_0 = \langle(1, a)\rangle, a \neq 0$$

$$V_0/1 = \{0\} = V_{e_1} \setminus 1, \text{ so } V_{e_1} = \langle(1, 0)\rangle = V_{ke_1} \text{ for } k \geq 1$$

$$\text{similarly } V_{ke_2} = \langle(0, 1)\rangle \text{ for } k \geq 1$$



# Example

3

$$(VF1) V_\alpha / i = V_{\alpha+e_i} \setminus i$$


$$(VF2) V_{\alpha+1} = \mathbf{1}V_\alpha$$

$$E = \{1, 2\}, V_0 = \langle(1, a)\rangle, a \neq 0$$

$$V_0/1 = \{0\} = V_{e_1} \setminus 1, \text{ so } V_{e_1} = \langle(1, 0)\rangle = V_{ke_1} \text{ for } k \geq 1$$

$$\text{similarly } V_{ke_2} = \langle(0, 1)\rangle \text{ for } k \geq 1$$

VF2 yields:





For  $W \subseteq K^E$  linear subspace,  $M(W)$  matroid on  $E$  defined by  $W$ .

For  $W \subseteq K^E$  linear subspace,  $M(W)$  matroid on  $E$  defined by  $W$ .

## **Theorem**

For a flock  $V$  on  $E$ , call  $I \subseteq E$  independent if  $\exists \alpha \in \mathbb{Z}^E$ :  $I$  independent in  $M(V_\alpha)$ . This defines a matroid  $M(V)$  on  $E$ .

For  $W \subseteq K^E$  linear subspace,  $M(W)$  matroid on  $E$  defined by  $W$ .

## Theorem

For a flock  $V$  on  $E$ , call  $I \subseteq E$  independent if  $\exists \alpha \in \mathbb{Z}^E$ :  $I$  independent in  $M(V_\alpha)$ . This defines a matroid  $M(V)$  on  $E$ .

## Question

Which matroids are flock-representable?

For  $W \subseteq K^E$  linear subspace,  $M(W)$  matroid on  $E$  defined by  $W$ .

## Theorem

For a flock  $V$  on  $E$ , call  $I \subseteq E$  independent if  $\exists \alpha \in \mathbb{Z}^E$ :  $I$  independent in  $M(V_\alpha)$ . This defines a matroid  $M(V)$  on  $E$ .

## Question

Which matroids are flock-representable?

## Theorem

Certainly all algebraic matroids over algebraically closed  $K$  of characteristic  $p > 0$  with  $\sigma(a) = a^{1/p}$ —so-called *Frobenius flocks*.

For  $W \subseteq K^E$  linear subspace,  $M(W)$  matroid on  $E$  defined by  $W$ .

## Theorem

For a flock  $V$  on  $E$ , call  $I \subseteq E$  independent if  $\exists \alpha \in \mathbb{Z}^E$ :  $I$  independent in  $M(V_\alpha)$ . This defines a matroid  $M(V)$  on  $E$ .

## Question

Which matroids are flock-representable?

## Theorem

Certainly all algebraic matroids over algebraically closed  $K$  of characteristic  $p > 0$  with  $\sigma(a) = a^{1/p}$ —so-called *Frobenius flocks*.

*But, (unfortunately?) many more.*

## Duality

If  $V$  is a flock over  $(K, \sigma)$ , then  $V^* : \alpha \mapsto V_{-\alpha}^\perp$  is a flock over  $(K, \sigma^{-1})$ , and it satisfies  $M(V^*) = M(V)^*$ .

## Duality

If  $V$  is a flock over  $(K, \sigma)$ , then  $V^* : \alpha \mapsto V_{-\alpha}^\perp$  is a flock over  $(K, \sigma^{-1})$ , and it satisfies  $M(V^*) = M(V)^*$ .

For a flock  $V$  on  $E$ ,  $\alpha \in \mathbb{Z}^E$ ,  $i \in E$ ,

- $i$  is a coloop in  $M(V_{\alpha+ke_i})$  for  $k \gg 0$  unless it is a loop in  $M(V)$ ;
- $i$  is a loop in  $M(V_{\alpha+ke_i})$  for  $k \ll 0$  unless it is a coloop in  $M(V)$ .

## Duality

If  $V$  is a flock over  $(K, \sigma)$ , then  $V^* : \alpha \mapsto V_{-\alpha}^\perp$  is a flock over  $(K, \sigma^{-1})$ , and it satisfies  $M(V^*) = M(V)^*$ .

For a flock  $V$  on  $E$ ,  $\alpha \in \mathbb{Z}^E$ ,  $i \in E$ ,

- $i$  is a coloop in  $M(V_{\alpha+ke_i})$  for  $k \gg 0$  unless it is a loop in  $M(V)$ ;
- $i$  is a loop in  $M(V_{\alpha+ke_i})$  for  $k \ll 0$  unless it is a coloop in  $M(V)$ .

## Minors

$V/i : \alpha \in \mathbb{Z}^{E-i} \rightarrow V_\beta/i$  for  $\beta|_{E-i} = \alpha$  and  $\beta_i \gg 0$ ; and

$V \setminus i : \alpha \in \mathbb{Z}^{E-i} \rightarrow V_\beta \setminus i$  for  $\beta|_{E-i} = \alpha$  and  $\beta_i \ll 0$

are flocks on  $E - i$ , satisfying

$M(V/i) = M(V)/i$ ,  $M(V \setminus i) = M(V) \setminus i$ , and  $V^*/i = (V \setminus i)^*$ .



$\sigma : K \rightarrow K$  trivial,  $W \subseteq K((t))^E$  a linear subspace  
for  $\alpha \in \mathbb{Z}^E$  set  $t^\alpha W := \{(t^{\alpha_i} w_i)_i \mid w \in W\} \subseteq K((t))^E$

$\sigma : K \rightarrow K$  trivial,  $W \subseteq K((t))^E$  a linear subspace  
for  $\alpha \in \mathbb{Z}^E$  set  $t^\alpha W := \{(t^{\alpha_i} w_i)_i \mid w \in W\} \subseteq K((t))^E$

## Construction

$V : \alpha \mapsto V_\alpha := \text{image in } K^E \text{ of } t^{-\alpha} W \cap K[[t]]^E$   
is a vector space flock of rank  $\dim W$  over  $(K, 1_K)$ .

$M(V_\alpha)$  has no loops if and only if  $\alpha \in \text{Trop}(W)$ .


*This suggests a tropical connection.*

$K$  algebraically closed,  $X \subseteq K^E$  irreducible variety  $\rightsquigarrow$  matroid  $M(X)$  on  $E$ :  $I$  independent iff  $X \rightarrow K^I$  dominant.

$K$  algebraically closed,  $X \subseteq K^E$  irreducible variety  $\rightsquigarrow$  matroid  $M(X)$  on  $E$ :  $I$  independent iff  $X \rightarrow K^I$  dominant.

## Ingleton's observation


If  $\text{char} K = 0$ , then  $M(X) = M(T_v X)$  for  $v \in X$  general, so  
 $\{\text{matroids algebraically representable over } K\}$   
 $= \{\text{matroids linearly representable over } K\}.$



$K$  algebraically closed,  $X \subseteq K^E$  irreducible variety  $\rightsquigarrow$  matroid  $M(X)$  on  $E$ :  $I$  independent iff  $X \rightarrow K^I$  dominant.

## Ingleton's observation


If  $\text{char} K = 0$ , then  $M(X) = M(T_v X)$  for  $v \in X$  general, so  
 $\{\text{matroids algebraically representable over } K\}$   
 $= \{\text{matroids linearly representable over } K\}.$



## Example

This is *not* true in char, say, 2:

$X = \{(x, y, z, xy, xz, yz, xzy) \mid x, y, z \in \overline{\mathbb{F}_2}^3\}$   
 represents the *non-Fano* matroid, which is  
 not linearly representable over  $\overline{\mathbb{F}_2}$ .



$K$  algebraically closed of char  $p$ ,  $\sigma : K \rightarrow K, a \mapsto a^{1/p}$   
 $\mathbb{Z}^E$  acts on  $K^E$  by homeomorphisms in the Zariski topology.

$K$  algebraically closed of char  $p$ ,  $\sigma : K \rightarrow K, a \mapsto a^{1/p}$   
 $\mathbb{Z}^E$  acts on  $K^E$  by homeomorphisms in the Zariski topology.

## Theorem

For  $X \subseteq K^E$  irreducible and  $v \in X$  general,  $V : \alpha \mapsto T_{\alpha v} \alpha X$  is a Frobenius flock over  $(K, \sigma)$  satisfying  $M(V) = M(X)$ .

$K$  algebraically closed of char  $p$ ,  $\sigma : K \rightarrow K, a \mapsto a^{1/p}$   
 $\mathbb{Z}^E$  acts on  $K^E$  by homeomorphisms in the Zariski topology.

## Theorem

For  $X \subseteq K^E$  irreducible and  $v \in X$  general,  $V : \alpha \mapsto T_{\alpha v} \alpha X$  is a Frobenius flock over  $(K, \sigma)$  satisfying  $M(V) = M(X)$ .

## Conditions (\*)

$v$  should satisfy the the following conditions *for all*  $\alpha \in \mathbb{Z}^E$ :

- $\alpha X$  is smooth at  $\alpha v$ , and
- $M(T_{\alpha v} \alpha X) = M(T_{\xi_\alpha} \alpha X)$ , where  $\xi$  is the generic point of  $\alpha X$ .

To reduce from *very general* to just *general* we establish finiteness properties of flocks.



# Example

$$X = \{(x, y, x + y, x + y^{(p^g)}) \mid (x, y) \in K^2\} \subseteq K^4, g > 1, M(X) = U_{2,4}$$

# Example

$$X = \{(x, y, x + y, x + y^{(p^g)}) \mid (x, y) \in K^2\} \subseteq K^4, g > 1, M(X) = U_{2,4}$$

$$T_0X = \text{row space of } \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}, \text{ so } 1, 4 \text{ parallel in } M(T_0X).$$

# Example

$$X = \{(x, y, x + y, x + y^{(p^g)}) \mid (x, y) \in K^2\} \subseteq K^4, g > 1, M(X) = U_{2,4}$$

$$T_0X = \text{row space of } \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}, \text{ so } 1, 4 \text{ parallel in } M(T_0X).$$

$$(-e_2 - e_3)X = \{(x, y, x^p + y, x + y^{(p^{g-1})}) \mid (x, y) \in K^2\}$$

$$T_0(-e_2 - e_3)X = \text{row space of } \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}; \text{ also } 2, 3 \text{ parallel.}$$

# Example

$$X = \{(x, y, x + y, x + y^{(p^g)}) \mid (x, y) \in K^2\} \subseteq K^4, g > 1, M(X) = U_{2,4}$$

$$T_0X = \text{row space of } \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}, \text{ so } 1, 4 \text{ parallel in } M(T_0X).$$


$$(-e_2 - e_3)X = \{(x, y, x^p + y, x + y^{(p^{g-1})}) \mid (x, y) \in K^2\}$$

$$T_0(-e_2 - e_3)X = \text{row space of } \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix}; \text{ also } 2, 3 \text{ parallel.}$$

$$(-ge_2 - ge_3)X = \{(x, y, x^{(p^g)} + y, x + y) \mid (x, y) \in K^2\}$$

$$T_0(-ge_2 - ge_3)X = \text{row space of } \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix}; 1, 4 \text{ indep.}$$

Cells where  $M(T_0(\alpha X)) = M(T_{\xi_\alpha} \alpha X)$  is constant:



These cells are *alcoved polytopes*: max-plus and min-plus closed.

## Definition (Dress-Wenzel)

A *matroid valuation* is a map  $\nu : \{d\text{-sets in } E\} \rightarrow \mathbb{R} \cup \{\infty\}$  such that  $\nu(B) \neq \infty$  for some  $B$  and  $\forall B, B', i \in B \setminus B' \exists j \in B' \setminus B :$   
 $\nu(B) + \nu(B') \geq \nu(B - i + j) + \nu(B' + i - j).$

( $\nu$  then lies in the *Dressian*)

## Definition (Dress-Wenzel)

A *matroid valuation* is a map  $\nu : \{d\text{-sets in } E\} \rightarrow \mathbb{R} \cup \{\infty\}$  such that  $\nu(B) \neq \infty$  for some  $B$  and  $\forall B, B', i \in B \setminus B' \exists j \in B' \setminus B : \nu(B) + \nu(B') \geq \nu(B - i + j) + \nu(B' + i - j)$ .

( $\nu$  then lies in the *Dressian*)

## Observations

$\nu \rightsquigarrow$  two matroids:  $M^\nu := \{B \mid \nu(B) < \infty\}$  and  $\{B \mid \nu(B) \text{ minimal}\}$ ; and  $\nu'(B) := \nu(B) - \alpha \cdot e_B$  is a valuation for each  $\alpha \in \mathbb{R}^E$ .

## Definition (Dress-Wenzel)

A *matroid valuation* is a map  $\nu : \{d\text{-sets in } E\} \rightarrow \mathbb{R} \cup \{\infty\}$  such that  $\nu(B) \neq \infty$  for some  $B$  and  $\forall B, B', i \in B \setminus B' \exists j \in B' \setminus B : \nu(B) + \nu(B') \geq \nu(B - i + j) + \nu(B' + i - j)$ .

( $\nu$  then lies in the *Dressian*)

## Observations

$\nu \rightsquigarrow$  two matroids:  $M^\nu := \{B \mid \nu(B) < \infty\}$  and  $\{B \mid \nu(B) \text{ minimal}\}$ ; and  $\nu'(B) := \nu(B) - \alpha \cdot e_B$  is a valuation for each  $\alpha \in \mathbb{R}^E$ .

## Theorem

Given a  $\mathbb{Z} \cup \{\infty\}$ -valued  $\nu$ , set  $M_\alpha^\nu := \{B \mid \nu(B) - \alpha \cdot e_B \text{ minimal}\}$  for each  $\alpha \in \mathbb{Z}^E$ . This satisfies matroid analogues of VF1, VF2. Conversely, each such *matroid flock* arises in this manner.



{algebraic varieties  $X \subseteq K^E$ }

$$(X, \nu) \mapsto (\alpha \mapsto T_{\alpha\nu}\alpha X)$$

$$\{\text{algebraic varieties } X \subseteq K^E\} \longrightarrow \{\text{Frobenius flocks } V : \alpha \mapsto V_\alpha\}$$

$$(X, \nu) \mapsto (\alpha \mapsto T_{\alpha\nu}\alpha X)$$

$$\{\text{algebraic varieties } X \subseteq K^E\} \longrightarrow \{\text{Frobenius flocks } V : \alpha \mapsto V_\alpha\}$$

$$V \mapsto (\alpha \mapsto M(V_\alpha))$$



$$\{\text{Matroid flocks } M : \alpha \mapsto M_\alpha\}$$

$$(X, \nu) \mapsto (\alpha \mapsto T_{\alpha\nu}\alpha X)$$

$$\{\text{algebraic varieties } X \subseteq K^E\} \longrightarrow \{\text{Frobenius flocks } V : \alpha \mapsto V_\alpha\}$$

$$V \mapsto (\alpha \mapsto M(V_\alpha))$$



$$\{\text{Matroid flocks } M : \alpha \mapsto M_\alpha\}$$

Murota—*thanks to Yu!*



$$\{\mathbb{Z} \cup \{\infty\}\text{-valued matroid valuations}\}$$

$$(X, \nu) \mapsto (\alpha \mapsto T_{\alpha\nu}\alpha X)$$

$$\{\text{algebraic varieties } X \subseteq K^E\} \longrightarrow \{\text{Frobenius flocks } V : \alpha \mapsto V_\alpha\}$$

$$X \mapsto M(X)$$

$$V \mapsto (\alpha \mapsto M(V_\alpha))$$

{Matroids on  $E$ }

{Matroid flocks  $M : \alpha \mapsto M_\alpha$ }

Murota—*thanks to Yu!*

$\{\mathbb{Z} \cup \{\infty\}$ -valued matroid valuations}

$$(X, \nu) \mapsto (\alpha \mapsto T_{\alpha\nu}\alpha X)$$

$$\{\text{algebraic varieties } X \subseteq K^E\} \longrightarrow \{\text{Frobenius flocks } V : \alpha \mapsto V_\alpha\}$$

$$X \mapsto M(X)$$

$$V \mapsto (\alpha \mapsto M(V_\alpha))$$

$$M \mapsto \bigcup_\alpha \{\text{bases of } M_\alpha\}$$

$$\{\text{Matroids on } E\} \longleftarrow \{\text{Matroid flocks } M : \alpha \mapsto M_\alpha\}$$

Murota—*thanks to Yu!*

$\{\mathbb{Z} \cup \{\infty\}$ -valued matroid valuations}

$$(X, \nu) \mapsto (\alpha \mapsto T_{\alpha\nu}\alpha X)$$

$$\{\text{algebraic varieties } X \subseteq K^E\} \longrightarrow \{\text{Frobenius flocks } V : \alpha \mapsto V_\alpha\}$$

$$X \mapsto M(X)$$

$$V \mapsto (\alpha \mapsto M(V_\alpha))$$

$$M \mapsto \bigcup_\alpha \{\text{bases of } M_\alpha\}$$

{Matroids on  $E$ }

{Matroid flocks  $M : \alpha \mapsto M_\alpha$ }


$$\nu \mapsto M^\nu$$

{ $\mathbb{Z} \cup \{\infty\}$ -valued matroid valuations}

Murota—*thanks to Yu!*

$$(X, \nu) \mapsto (\alpha \mapsto T_{\alpha\nu}\alpha X)$$

$$\{\text{algebraic varieties } X \subseteq K^E\} \longrightarrow \{\text{Frobenius flocks } V : \alpha \mapsto V_\alpha\}$$



So to a  $d$ -dimensional algebraic variety  $X \subseteq K^E$  in char  $p$  we associate the *Lindstrom valuation*  $\nu^X : \{d\text{-subsets of } E\} \rightarrow \mathbb{Z} \cup \{\infty\}$ .

**Cartwright found a direct construction of  $\nu^X$ .**



## Proposition

$\nu$  a  $\mathbb{Z} \cup \{\infty\}$ -valued valuation and  $\alpha, \beta \in \mathbb{R}^E$ . Then  $M_\alpha^\nu \supseteq M_\beta^\nu$  iff  $\forall i \neq j \forall B \in M_\beta^\nu : \alpha_i - \alpha_j \geq \nu(B) - \nu(B - i + j)$ .

This gives a finite polyhedral complex with *alcoved polyhedra*, in the relative interior of which  $M_\alpha^\nu$  is constant.

## Proposition

$\nu$  a  $\mathbb{Z} \cup \{\infty\}$ -valued valuation and  $\alpha, \beta \in \mathbb{R}^E$ . Then  $M_\alpha^\nu \supseteq M_\beta^\nu$  iff  $\forall i \neq j \forall B \in M_\beta^\nu : \alpha_i - \alpha_j \geq \nu(B) - \nu(B - i + j)$ .

This gives a finite polyhedral complex with *alcoved polyhedra*, in the relative interior of which  $M_\alpha^\nu$  is constant.

## Consequences:

- A vector space flock can be specified by a finite amount of data.
- Conditions (\*) on  $\nu$  are satisfied by general  $\nu$ : if  $\alpha\nu$  satisfies it for  $\alpha X$ , and if  $M_\alpha^{\nu_X} = M_{\alpha+e_J}^{\nu_X}$ , then  $(e_J + \alpha)\nu$  satisfies it at  $(\alpha + e_J)X$ .

## **Definition (Dress-Wenzel)**

A matroid  $M$  is *rigid* if every valuation  $\nu$  with  $M^\nu = M$  is of the form  $M \rightarrow \mathbb{R}$ ,  $B \mapsto \alpha \cdot e_B$  for some  $\alpha \in \mathbb{R}^E$ .

## Definition (Dress-Wenzel)

A matroid  $M$  is *rigid* if every valuation  $\nu$  with  $M^\nu = M$  is of the form  $M \rightarrow \mathbb{R}$ ,  $B \mapsto \alpha \cdot e_B$  for some  $\alpha \in \mathbb{R}^E$ .

## Theorem

A rigid matroid is algebraically representable over an algebraically closed field  $K$  of positive characteristic if and only if it is linearly representable over  $K$ .

## Definition (Dress-Wenzel)

A matroid  $M$  is *rigid* if every valuation  $\nu$  with  $M^\nu = M$  is of the form  $M \rightarrow \mathbb{R}$ ,  $B \mapsto \alpha \cdot e_B$  for some  $\alpha \in \mathbb{R}^E$ .

## Theorem

A rigid matroid is algebraically representable over an algebraically closed field  $K$  of positive characteristic if and only if it is linearly representable over  $K$ .

## Proof

If  $X$  is an algebraic representation, then the Lindström valuation  $\nu^X : M(X) \rightarrow \mathbb{Z}$  sends  $B \mapsto \alpha \cdot e_B$  for some  $\alpha \in \mathbb{R}^E$ . Then  $M_\alpha^\nu = M^\nu$ , and the cell where this happens also contains an integral  $\alpha$ . Now  $M(X) = M(T_{\alpha\nu}\alpha X)$  for  $\nu \in X$  general.  $\square$

Using this theorem, we can re-prove several existing results, such as: the projective plane over  $\mathbb{F}_p$  is algebraically representable only over fields of characteristic  $p$  (Lindström).

To extend the applicability of this method, we are trying to relax the condition that the matroid be rigid.

Using this theorem, we can re-prove several existing results, such as: the projective plane over  $\mathbb{F}_p$  is algebraically representable only over fields of characteristic  $p$  (Lindström).


To extend the applicability of this method, we are trying to relax the condition that the matroid be rigid.

*However*, using Frobenius flocks alone one cannot reprove all non-algebraicity results:

## **Theorem-in-progress**

If  $M$  has a flock representation over  $(K, \sigma)$ , and  $H$  is a circuit hyperplane in  $M$ , then the matroid obtained by turning  $H$  into a basis again has a flock representation over  $(K, \sigma)$ .


The missing diagonal plane is independent:



Vamos has a flock realisation; but it is non-algebraic.



The missing diagonal plane is independent:




Vamos has a flock realisation; but it is non-algebraic.

## Questions/projects

- Compute the Lindström valuation from a prime ideal? (Bollen)
- Bounds on the Lindström locus in the Dressian? (No idea.)
- ...

The missing diagonal plane is independent:



Vamos has a flock realisation; but it is non-algebraic.

## Questions/projects

- Compute the Lindström valuation from a prime ideal? (Bollen)
- Bounds on the Lindström locus in the Dressian? (No idea.)
- ...

*Thanks!*